

DR.K.S.CHAUDHARI
M.S.[ENT].D.O.R.L.M.B.B.S.FICCS[COSMETIC]
COSMETIC SURGEON
Cosmetic Laser Surgery Center
Prasad chambers,27&24
Near Excel petrol pump
Karve road,Pune-411004.
Clinic-020-25441104/Res. 020-25432118
Mobile.98220-58916
kchaudhari@vsnl.com
www.cosmeticsurgerypune.com
Time:10 to 1pm.6to 8pm
Sunday closed

CONSENT FORM FOR LIPOSUCTION

INSTRUCTIONS

This is an informed-consent document that has been prepared to help inform you concerning suction-assisted liposuction (“liposuction”) surgery, its risks, and alternative treatment.

It is important that you read this information carefully and completely. Please initial each page, indicating that you have read the page and sign the consent for surgery as proposed by Dr.K.S.Chaudhari

INTRODUCTION

Suction-assisted liposuction is a surgical technique to remove unwanted deposits of fat from specific areas of the body, including the face and neck, upper arms, trunk, abdomen, buttocks, hips and thighs, and the knees, calves and ankles. Liposuction is not a substitute for weight reduction, but a method for removing localized deposits of fatty tissue that does not respond to diet or exercise. Suction-assisted liposuction may be performed as a primary procedure for body contouring or combined with other surgical techniques such as facelift, abdominoplasty, or thigh lift procedures to tighten loose skin and supporting structures.

The best candidates for liposuction are individuals of relatively normal weight who have excess fat in particular body areas. Having firm, elastic skin will result in a better final contour after liposuction. Hanging skin will not reshape itself to the new contours and may require additional surgical techniques to remove and tighten excess skin. Body-contour irregularities due to structures other than fat cannot be improved by suction-liposuction. Suction-assisted liposuction by itself may not improve areas of dimpled skin known as “cellulite.”

ALTERNATIVE TREATMENT

Alternative forms of management consist of not treating the areas of fatty deposits. Diet and exercise regimens may be of benefit in the overall reduction of excess body fat. Direct removal of excess skin and fatty tissue may be necessary in addition to suction-assisted liposuction in some patients.

Risks and potential complications are associated with alternative forms of treatment that involve surgery.

RISKS of SUCTION-ASSISTED LIPOSUCTION SURGERY

Every surgical procedure involves a certain amount of risk, and it is important that you understand the risks involved with suction-assisted liposuction. An individual's choice to undergo a surgical procedure is based on the comparison of the risk to potential benefit. Although the majority of patients do not experience these complications, you should discuss each of them with your surgeon to make sure you understand the risks, potential complications, and consequences of suction-assisted liposuction.

Patient selection- Individuals with poor skin tone, medical problems, obesity, or unrealistic expectations may not be candidates for suction-assisted liposuction.

Bleeding- It is possible, though unusual, to have a bleeding episode during or after surgery. Should post-operative bleeding occur, it may require emergency treatment to drain accumulated blood or blood transfusion. Do not take any aspirin or anti-inflammatory medications for three weeks before surgery, as this may increase the risk of bleeding.

Infection- An infection is quite unusual after this type of surgery. Should an infection occur, treatment including antibiotics or additional surgery may be necessary.

Change in and skin sensation- Temporary changes in skin sensation after suction-assisted liposuction occur which usually resolve. Diminished (or complete loss of skin sensation) infrequently occurs and may not completely resolve.

Skin scarring- Although good wound healing after a surgical procedure is expected, abnormal scars may occur within the skin and deeper tissues. abnormal scars may result. Scars may be unattractive and of different color than surrounding skin. Additional treatments including surgery may be needed to treat abnormal scarring.

Skin contour irregularities- Contour irregularities and depressions in the skin may occur after suction-assisted liposuction. Visible and palpable wrinkling of skin can occur. Additional treatments including surgery may be necessary to treat skin contour irregularities following suction-assisted liposuction.

Asymmetry- Symmetrical body appearance may not result from suction-assisted liposuction surgery. Factors such as skin tone, bony prominence, and muscle tone may contribute to normal asymmetry in body features.

Surgical shock- In rare circumstances, this procedure can cause severe trauma, particularly when multiple or extensive areas are suctioned at one time. Although serious complications are infrequent, infections or excessive fluid loss can lead to severe illness and even death. Should surgical shock occur after suction-assisted liposuction, hospitalization and additional treatment would be necessary.

Pulmonary complications- Fat embolism syndrome occurs when fat droplets are trapped in the lungs. This is a very rare and possibly fatal complication of suction-assisted liposuction. Should fat embolism or other pulmonary

complications occur following suction-assisted liposuction, additional treatment including hospitalization may be necessary.

Skin loss- Skin loss is rare after suction-assisted liposuction. Additional treatments including surgery may be necessary.

Seroma- Fluid accumulations infrequently occur in areas where suction-assisted liposuction has been performed. Additional treatments or surgery to drain accumulations of fluid may be necessary.

Long term effects- Subsequent alterations in body contour may occur as the result of aging, weight loss or gain, pregnancy, or other circumstances not related to suction-assisted liposuction.

Allergic reactions- In rare cases, local allergies to tape, suture material, or topical preparations have been reported.

Systemic reactions which are more serious may occur to drugs used during surgery and prescription medicines. Allergic reactions may require additional treatment.

Other- You may be disappointed with the results of surgery. It may be necessary to perform additional surgery to improve your results.

Surgical anesthesia- Both local and general anesthesia involves risk. There is the possibility of complications, injury, and even death from all forms of surgical anesthesia or sedation.

I _____ have read above mentioned information & have understood it fully. I permit Dr.K.S.Chaudhari To perform Liposuction on my-----
Under local anesthesia & also permit for medical photography before during & after surgery for medical records & research.

Patient signature.

Date

Witness